



UCB and Dermira Announce U.S. and EU Regulatory Submissions for CIMZIA® (certolizumab pegol) for the treatment of Moderate-to-Severe Chronic Plaque Psoriasis

- *The regulatory submissions for this new indication seek to expand the use of CIMZIA for the treatment of patients with moderate-to-severe chronic plaque psoriasis*
- *Data from three positive Phase 3 studies support the regulatory submissions*

Brussels, Belgium and Menlo Park, Calif., July 25, 2017 — UCB (Euronext: UCB) and Dermira, Inc. (NASDAQ: DERM) today announced the UCB submission of a supplemental Biologics License Application (sBLA) to the U.S. Food and Drug Administration (FDA) for CIMZIA® (certolizumab pegol). Separately, UCB also submitted a regulatory filing with the European Medicines Agency (EMA). Both regulatory filings seek to expand the approved indications for CIMZIA to include treatment of adult patients with moderate-to-severe chronic plaque psoriasis. An additional submission to expand the use of CIMZIA in this patient population is also planned with Health Canada.

The Phase 3 CIMZIA clinical development program in psoriasis was led by Dermira, Inc., in collaboration with UCB as the regulatory sponsor. Under the terms of the agreement announced in July 2014, Dermira obtained exclusive rights to develop CIMZIA in psoriasis in the United States, Canada and the EU. Subject to regulatory approval of CIMZIA in psoriasis, Dermira is granted an exclusive commercial license to market CIMZIA to dermatologists in the United States and Canada. UCB will retain marketing rights for CIMZIA in the U.S. and Canada for rheumatoid arthritis, psoriatic arthritis, Crohn's disease and ankylosing spondylitis. In the rest of world, UCB retains marketing rights for CIMZIA for all approved indications, including psoriasis, subject to approval.

“UCB is proud to have achieved this important regulatory milestone in partnership with Dermira. Based on the results of our Phase 3 clinical trial program, we believe CIMZIA, the only Fc-free, PEGylated anti-TNF, has the potential to be a valuable treatment option for patients living with moderate-to-severe chronic plaque psoriasis. This underserved patient population faces a heavy disease burden, including pain and a reduced quality of life. Our psoriasis development program, and our collaboration with Dermira, aims to provide value and broaden access to CIMZIA,” said Emmanuel Caeymaex, Head of Immunology and Executive Vice President, Immunology Patient Value Unit, UCB

“Despite the tremendous progress that has been made in psoriasis drug development during the last decade and the number of FDA-approved treatment options that are currently available, healthcare professionals and patients continue to express the need for additional treatment options,” said Tom Wiggins, chairman and chief executive officer of Dermira. “If approved, we believe CIMZIA would represent an important new treatment option for previously untreated patients living with moderate-to-severe chronic plaque psoriasis, as well as patients who have received prior treatment with another biologic therapy.”

The regulatory submissions are based on results from three Phase 3 clinical trials – CIMPASI-1, CIMPASI-2 and CIMPACT, which each evaluated the efficacy and safety of CIMZIA in adult patients with moderate-to-severe chronic plaque psoriasis. The co-primary endpoints evaluated in both CIMPASI-1 and CIMPASI-2 were the percentage of patients who achieved a 75% or greater disease improvement from baseline as measured by the Psoriasis Area and Severity Index (PASI 75) and the percentage of patients achieving at least a two-point improvement on a five-point Physician’s Global Assessment (PGA) scale to a final score representing clear or almost clear skin, each compared with placebo, at week 16. The primary endpoint in CIMPACT was the percentage of patients on CIMZIA achieving a PASI 75 response, compared with placebo, at week 12. Secondary endpoints of the CIMPACT trial included a comparison of the efficacy of CIMZIA to ENBREL® (etanercept) as measured by PASI 75 response at week 12 and the percentage of patients who achieved at least a two-point improvement to a final score representing clear or almost clear skin on the five-point PGA scale, at week 12.

The three Phase 3 [trials](#) enrolled approximately 1,000 patients, including patients with and without prior treatment experience with biologic products. In CIMPASI-1 and CIMPASI-2, patients were randomized to one of three dosing arms—400 mg every two weeks, 400 mg at weeks 0, 2, and 4 followed by 200 mg every two weeks, or placebo every two weeks. In CIMPACT, patients were randomized to one of four dosing arms—CIMZIA at 400 mg every two weeks, CIMZIA at 400 mg at weeks 0, 2, and 4 followed by 200 mg every two weeks, ENBREL at 50 mg twice weekly, or placebo every two weeks. In all three trials, CIMZIA demonstrated statistically significant improvements for all primary or co-primary endpoints compared to placebo at all treatment doses.

CIMZIA is not currently approved for the treatment of psoriasis by any regulatory authority worldwide.

About Psoriasis

Psoriasis is a common, chronic, immune-mediated inflammatory disorder with primary involvement of the skin. It affects nearly three percent of the population, or approximately 125 million people worldwide. The skin condition affects men and women of all ages and ethnicities. Psoriasis signs and symptoms can vary, but may include red patches of skin covered with silvery scales, dry, cracked skin that may bleed and thickened, pitted or ridged nails.²

About Cimzia® in the U.S.

Cimzia® is the only Fc-free, PEGylated anti-TNF (Tumor Necrosis Factor). Cimzia® has a high affinity for human TNF-alpha, selectively neutralizing the pathophysiological effects of TNF-alpha.

Cimzia® is indicated for the treatment of adults with moderately to severely active rheumatoid arthritis, adults with active psoriatic arthritis (PsA), and adults with active ankylosing spondylitis (AS). In addition, it is indicated for reducing signs and symptoms of Crohn's disease and maintaining clinical response in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. See important safety information including risk of serious bacterial, viral and fungal infections and tuberculosis below.

Important Safety Information about Cimzia® in the United States

Risk of Serious Infections and Malignancy

Patients treated with Cimzia® are at an increased risk for developing serious infections that may lead to hospitalization or death. Most patients who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids. Cimzia® should be discontinued if a patient develops a serious infection or sepsis. Reported infections include:

- **Active tuberculosis, including reactivation of latent tuberculosis.** Patients with tuberculosis have frequently presented with disseminated or extrapulmonary disease. Patients should be tested for latent tuberculosis before Cimzia® use and during therapy. Treatment for latent infection should be initiated prior to Cimzia® use.
- **Invasive fungal infections, including histoplasmosis, coccidioidomycosis, candidiasis, aspergillosis, blastomycosis, and pneumocystosis.** Patients with histoplasmosis or other invasive fungal infections may present with disseminated, rather than localized disease. Antigen and antibody testing for histoplasmosis may be negative in some patients with active infection. Empiric anti-fungal therapy should be considered in patients at risk for invasive fungal infections who develop severe systemic illness.
- **Bacterial, viral and other infections due to opportunistic pathogens, including Legionella and Listeria.**

The risks and benefits of treatment with Cimzia® should be carefully considered prior to initiating therapy in patients with chronic or recurrent infection. Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with Cimzia®, including the possible development of tuberculosis in patients who tested negative for latent tuberculosis infection prior to initiating therapy.

Lymphoma and other malignancies, some fatal, have been reported in children and adolescent patients treated with TNF blockers, of which Cimzia® is a member. Cimzia® is not indicated for use in pediatric patients.

Patients treated with Cimzia® are at an increased risk for developing serious infections involving various organ systems and sites that may lead to hospitalization or death. Opportunistic infections due to bacterial, mycobacterial, invasive fungal, viral, parasitic, or other opportunistic pathogens including aspergillosis, blastomycosis, candidiasis, coccidioidomycosis, histoplasmosis, legionellosis, listeriosis, pneumocystosis and tuberculosis have been reported with TNF blockers. Patients have frequently presented with disseminated rather than localized disease.

Treatment with Cimzia® should not be initiated in patients with an active infection, including clinically important localized infections. Cimzia® should be discontinued if a patient develops a serious infection or sepsis. Patients greater than 65 years of age, patients with co-morbid conditions, and/or patients taking concomitant immunosuppressants (e.g., corticosteroids or methotrexate) may be at a greater risk of infection. Patients who develop a new infection during treatment with Cimzia® should be closely monitored, undergo a prompt and complete diagnostic workup appropriate for immunocompromised patients, and appropriate

antimicrobial therapy should be initiated. Appropriate empiric antifungal therapy should also be considered while a diagnostic workup is performed for patients who develop a serious systemic illness and reside or travel in regions where mycoses are endemic.

Malignancies

During controlled and open-labeled portions of Cimzia® studies of Crohn's disease and other diseases, malignancies (excluding non-melanoma skin cancer) were observed at a rate of 0.5 per 100 patient-years among 4,650 Cimzia®-treated patients versus a rate of 0.6 per 100 patient-years among 1,319 placebo-treated patients. In studies of Cimzia® for Crohn's disease and other investigational uses, there was one case of lymphoma among 2,657 Cimzia®-treated patients and one case of Hodgkin lymphoma among 1,319 placebo-treated patients. In Cimzia® RA clinical trials (placebo-controlled and open label), a total of three cases of lymphoma were observed among 2,367 patients. This is approximately 2-fold higher than expected in the general population. Patients with RA, particularly those with highly active disease, are at a higher risk for the development of lymphoma. The potential role of TNF blocker therapy in the development of malignancies is not known.

Malignancies, some fatal, have been reported among children, adolescents, and young adults who received treatment with TNF-blocking agents (initiation of therapy ≤ 18 years of age), of which Cimzia® is a member. Approximately half of the cases were lymphoma (including Hodgkin's and non-Hodgkin's lymphoma), while the other cases represented a variety of different malignancies and included rare malignancies associated with immunosuppression and malignancies not usually observed in children and adolescents. Most of the patients were receiving concomitant immunosuppressants.

Cases of acute and chronic leukemia have been reported with TNF-blocker use. Even in the absence of TNF-blocker therapy, patients with RA may be at a higher risk (approximately 2-fold) than the general population for developing leukemia.

Postmarketing cases of hepatosplenic T-cell lymphoma (HSTCL), a rare type of T-cell lymphoma that has a very aggressive disease course and is usually fatal, have been reported in patients treated with TNF blockers, including Cimzia®. The majority of reported TNF blocker cases occurred in adolescent and young adult males with Crohn's disease or ulcerative colitis. Almost all of these patients had received treatment with the immunosuppressants azathioprine and/or 6-mercaptopurine (6-MP) concomitantly with a TNF blocker at or prior to diagnosis. Carefully assess the risks and benefits of treatment with Cimzia®, especially in these patient types.

Melanoma and Merkel cell carcinoma have been reported in patients treated with TNF-antagonists, including Cimzia®. Periodic skin examinations are recommended for all patients, particularly those with risk factors for skin cancer.

Heart Failure

Cases of worsening congestive heart failure (CHF) and new onset CHF have been reported with TNF blockers. Cimzia® has not been formally studied in patients with CHF. Exercise caution when using Cimzia® in patients who have heart failure and monitor them carefully.

Hypersensitivity

Symptoms compatible with hypersensitivity reactions, including angioedema, dyspnea, hypotension, rash, serum sickness, and urticaria, have been reported rarely following Cimzia® administration. Some of these reactions occurred after the first administration of Cimzia®. If such reactions occur, discontinue further administration of Cimzia® and institute appropriate therapy.

Hepatitis B Reactivation

Use of TNF blockers, including Cimzia®, has been associated with reactivation of hepatitis B virus (HBV) in patients who are chronic carriers of this virus. Some cases have been fatal. Test patients for HBV infection before initiating treatment with Cimzia®. Exercise caution in prescribing Cimzia® for patients identified as carriers of HBV, with careful evaluation and monitoring prior to and during treatment. In patients who develop HBV reactivation, discontinue Cimzia® and initiate effective anti-viral therapy with appropriate supportive treatment.

Neurologic Reactions

Use of TNF blockers, including Cimzia®, has been associated with rare cases of new onset or exacerbation of clinical symptoms and/or radiographic evidence of central nervous system demyelinating disease, including multiple sclerosis, and with peripheral demyelinating disease, including Guillain-Barré syndrome. Rare cases of neurological disorders, including seizure disorder, optic neuritis, and peripheral neuropathy have been reported in patients treated with Cimzia®. Exercise caution in considering the use of Cimzia® in patients with these disorders.

Hematologic Reactions

Rare reports of pancytopenia, including aplastic anemia, have been reported with TNF blockers. Medically significant cytopenia (e.g., leukopenia, pancytopenia, thrombocytopenia) has been infrequently reported with Cimzia®. Advise all patients to seek immediate medical attention if they develop signs and symptoms suggestive of blood dyscrasias or infection (e.g., persistent fever, bruising, bleeding, pallor) while on Cimzia®. Consider discontinuation of Cimzia® therapy in patients with confirmed significant hematologic abnormalities.

Drug Interactions

An increased risk of serious infections has been seen in clinical trials of other TNF blocking agents used in combination with anakinra or abatacept. Formal drug interaction studies have not been performed with rituximab or natalizumab; however, because of the nature of the adverse events seen with these combinations with TNF blocker therapy, similar toxicities may also result from the use of Cimzia® in these combinations. Therefore, the combination of Cimzia® with anakinra, abatacept, rituximab, or natalizumab is not recommended. Interference with certain coagulation assays has been detected in patients treated with Cimzia®. There is no evidence that Cimzia® therapy has an effect on in vivo coagulation. Cimzia® may cause erroneously elevated aPTT assay results in patients without coagulation abnormalities.

Autoimmunity

Treatment with Cimzia® may result in the formation of autoantibodies and, rarely, in the development of a lupus-like syndrome. Discontinue treatment if symptoms of lupus-like syndrome develop.

Immunizations

Do not administer live vaccines or live-attenuated vaccines concurrently with Cimzia®.

Adverse Reactions

In controlled Crohn's clinical trials, the most common adverse events that occurred in ≥5% of Cimzia® patients (n=620) and more frequently than with placebo (n=614) were upper respiratory infection (20% Cimzia®, 13% placebo), urinary tract infection (7% Cimzia®, 6% placebo), and arthralgia (6% Cimzia®, 4% placebo). The proportion of patients who discontinued treatment due to adverse reactions in the controlled clinical studies was 8% for Cimzia® and 7% for placebo.

In controlled RA clinical trials, the most common adverse events that occurred in ≥3% of patients taking Cimzia® 200 mg every other week with concomitant methotrexate (n=640) and more frequently than with placebo with concomitant methotrexate (n=324) were upper respiratory tract infection (6% Cimzia®, 2% placebo), headache (5% Cimzia®, 4% placebo), hypertension (5% Cimzia®, 2% placebo), nasopharyngitis (5% Cimzia®, 1% placebo), back pain (4% Cimzia®, 1% placebo), pyrexia (3% Cimzia®, 2% placebo), pharyngitis (3% Cimzia®, 1% placebo), rash (3% Cimzia®, 1% placebo), acute bronchitis (3% Cimzia®, 1% placebo), fatigue (3% Cimzia®, 2% placebo). Hypertensive adverse reactions were observed more frequently in patients receiving Cimzia® than in controls. These adverse reactions occurred more frequently among patients with a baseline history of hypertension and among patients receiving concomitant corticosteroids and non-steroidal anti-inflammatory drugs. Patients receiving Cimzia® 400 mg as monotherapy every 4 weeks in RA controlled clinical trials had similar adverse reactions to those patients receiving Cimzia® 200 mg every other week. The proportion of patients who discontinued treatment due to adverse reactions in the controlled clinical studies was 5% for Cimzia® and 2.5% for placebo.

The safety profile for patients with Psoriatic Arthritis (PsA) treated with CIMZIA® was similar to the safety profile seen in patients with RA and previous experience with Cimzia®.

The safety profile for AS patients treated with Cimzia® was similar to the safety profile seen in patients with RA.

For full prescribing information, please visit www.ucb.com

CIMZIA® is a registered trademark of the UCB Group of Companies.

About Cimzia® in the EU/EEA

In the EU, Cimzia® in combination with methotrexate (MTX) is indicated for the treatment of moderate to severe active RA in adult patients inadequately responsive to disease-modifying anti-rheumatic drugs (DMARDs) including MTX.

Cimzia® can be given as monotherapy in case of intolerance to MTX or when continued treatment with MTX is inappropriate. CIMZIA® in combination with MTX is also indicated for the treatment of severe, active and progressive RA in adults not previously treated with MTX or other DMARDs.

Cimzia® has been shown to reduce the rate of progression of joint damage as measured by X-ray and to improve physical function, when given in combination with MTX.

Cimzia®, in combination with MTX, is also indicated for the treatment of active psoriatic arthritis in adults when the response to previous DMARD therapy has been inadequate. Cimzia® can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate.

Cimzia® is also indicated in the EU for the treatment of adult patients with severe active axial spondyloarthritis (axSpA), comprising:

- Ankylosing spondylitis (AS) - adults with severe active AS who have had an inadequate response to, or are intolerant to non-steroidal anti-inflammatory drugs (NSAIDs).
- Axial spondyloarthritis (axSpA) without radiographic evidence of AS - adults with severe active axSpA without radiographic evidence of AS but with objective signs of inflammation by elevated C-reactive protein (CRP) and/or Magnetic Resonance Imaging (MRI) who have had an inadequate response to, or are intolerant to NSAIDs.⁴

Important Safety Information about Cimzia® in the EU/EEA

Cimzia® was studied in 4,049 patients with rheumatoid arthritis (RA) in controlled and open label trials for up to 92 months. The commonly reported adverse reactions (1-10%) in clinical trials with Cimzia® and post-marketing were viral infections (includes herpes, papillomavirus, influenza), bacterial infections (including abscess), rash, headache (including migraine), asthenia, leukopenia (including lymphopenia, neutropenia), eosinophilic disorder, pain (any sites), pyrexia, sensory abnormalities, hypertension, pruritus (any sites), hepatitis (including hepatic enzyme increase), injection site reactions, and nausea. Serious adverse reactions include sepsis, opportunistic infections, tuberculosis, herpes zoster, lymphoma, leukaemia, solid organ tumours, angioneurotic oedema, cardiomyopathies (includes heart failure), ischemic coronary artery disorders, pancytopenia, hypercoagulation (including thrombophlebitis, pulmonary embolism), cerebrovascular accident, vasculitis, hepatitis/hepatopathy (includes cirrhosis), and renal impairment/nephropathy (includes nephritis). In RA controlled clinical trials, 4.4% of patients discontinued taking Cimzia® due to adverse events vs. 2.7% for placebo.

Cimzia® is contraindicated in patients with hypersensitivity to the active substance or any of the excipients, active tuberculosis or other severe infections such as sepsis or opportunistic infections or moderate-to-severe heart failure.

Serious infections including sepsis, tuberculosis and opportunistic infections have been reported in patients receiving Cimzia®. Some of these events have been fatal. Monitor patients closely for signs and symptoms of infections including tuberculosis before, during and after treatment with Cimzia®. Treatment with Cimzia® must not be initiated in patients with a clinically important active infection. If an infection develops, monitor

carefully and stop Cimzia® if infection becomes serious. Before initiation of therapy with Cimzia®, all patients must be evaluated for both active and inactive (latent) tuberculosis infection. If active tuberculosis is diagnosed prior to or during treatment, Cimzia® therapy must not be initiated and must be discontinued. If latent tuberculosis is diagnosed, appropriate anti-tuberculosis therapy must be started before initiating treatment with Cimzia®. Patients should be instructed to seek medical advice if signs/symptoms (e.g. persistent cough, wasting/weight loss, low grade fever, listlessness) suggestive of tuberculosis occur during or after therapy with Cimzia®.

Reactivation of hepatitis B has occurred in patients receiving a TNF-antagonist including Cimzia® who are chronic carriers of the virus (i.e. surface antigen positive). Some cases have had a fatal outcome. Patients should be tested for HBV infection before initiating treatment with Cimzia®. Carriers of HBV who require treatment with Cimzia® should be closely monitored and in the case of HBV reactivation Cimzia® should be stopped and effective anti-viral therapy with appropriate supportive treatment should be initiated.

TNF antagonists including Cimzia® may increase the risk of new onset or exacerbation of clinical symptoms and/or radiographic evidence of demyelinating disease; of formation of autoantibodies and uncommonly of the development of a lupus-like syndrome; of severe hypersensitivity reactions. If a patient develops any of these adverse reactions, Cimzia® should be discontinued and appropriate therapy instituted.

With the current knowledge, a possible risk for the development of lymphomas, leukaemia or other malignancies in patients treated with a TNF antagonist cannot be excluded. Rare cases of neurological disorders, including seizure disorder, neuritis and peripheral neuropathy, have been reported in patients treated with Cimzia®.

Adverse reactions of the hematologic system, including medically significant cytopaenia, have been infrequently reported with Cimzia®. Advise all patients to seek immediate medical attention if they develop signs and symptoms suggestive of blood dyscrasias or infection (e.g., persistent fever, bruising, bleeding, pallor) while on Cimzia®. Consider discontinuation of Cimzia® therapy in patients with confirmed significant haematological abnormalities.

The use of Cimzia® in combination with anakinra or abatacept is not recommended due to a potential increased risk of serious infections. As no data are available, Cimzia® should not be administered concurrently with live vaccines. The 14-day half-life of Cimzia® should be taken into consideration if a surgical procedure is planned. A patient who requires surgery while on CIMZIA® should be closely monitored for infections.

Cimzia® was studied in 325 patients with active axial spondyloarthritis (axSpA) in a placebo-controlled clinical trial for up to 30 months and in 409 patients with psoriatic arthritis (PsA) in a placebo-controlled clinical trial for up to 30 months. The safety profile for axSpA and PsA patients treated with Cimzia® was consistent with the safety profile in RA and previous experience with Cimzia®.

Please consult the full prescribing information in relation to other side effects, full safety and prescribing information. European SmPC date of revision 18th May 2017.

http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/001037/WC500069763.pdf

References

1. UCB Data on File.
2. International Federation of Psoriasis Associations. Accessed July 14, 2016 at <http://www.worldpsoriasisday.com/web/page.aspx?refid=130>

CIMZIA® is a registered trademark of the UCB Group of Companies.

ENBREL® (etanercept) is a registered trademark of Amgen Inc.

About Dermira

Dermira is a biopharmaceutical company dedicated to bringing biotech ingenuity to medical dermatology by delivering differentiated, new therapies to the millions of patients living with chronic skin conditions. Dermira is committed to understanding the needs of both patients and physicians and using its insight to identify and develop leading-edge medical dermatology programs. Dermira's product pipeline includes three late-stage product candidates that could have a profound impact on the lives of patients: glycopyrronium tosylate (formerly DRM04), which has completed a Phase 3 program for the treatment of primary axillary hyperhidrosis (excessive underarm sweating); CIMZIA® (certolizumab pegol), in Phase 3 development in collaboration with UCB Pharma S.A. for the treatment of moderate-to-severe chronic plaque psoriasis; and olumacostat glasaretil (formerly DRM01), in Phase 3 development for the treatment of acne vulgaris. Dermira is headquartered in Menlo Park, Calif. For more information, please visit www.dermira.com.

In addition to filings with the Securities and Exchange Commission (SEC), press releases, public conference calls and webcasts, Dermira uses its website (www.dermira.com) and LinkedIn page (<https://www.linkedin.com/company/dermira-inc>) as channels of distribution of information about its company, product candidates, planned financial and other announcements, attendance at upcoming investor and industry conferences and other matters. Such information may be deemed material information and Dermira may use these channels to comply with its disclosure obligations under Regulation FD. Therefore, investors should monitor Dermira's website and LinkedIn page in addition to following its SEC filings, press releases, public conference calls and webcasts.

About UCB

UCB, Brussels, Belgium (www.ucb.com) is a global biopharmaceutical company focused on the discovery and development of innovative medicines and solutions to transform the lives of people living with severe diseases of the immune system or of the central nervous system. With more than 7,500 people in approximately 40 countries, the company generated revenue of € 4.2 billion in 2016. UCB is listed on Euronext Brussels (symbol: UCB). Follow us on Twitter: @UCB_news.

Dermira Forward-Looking Statements

The information in this press release contains forward-looking statements and information within the meaning of Section 27A of the Securities Act of 1933, as amended, and Section 21E of the Securities Exchange Act of 1934, as amended, which are subject to the "safe harbor" created by those sections. This press release contains forward-looking statements that involve substantial risks and uncertainties, including statements with respect to CIMZIA becoming an important treatment option for, and providing value to, patients living

with psoriasis; the broadening of access to CIMZIA; anticipated submission of marketing applications to regulatory authorities in the U.S., Canada and EU to support potential approvals for CIMZIA as a treatment option for patients with moderate-to-severe chronic plaque psoriasis and timing expectations for such submission; and potential approval of CIMZIA as a treatment option for moderate-to-severe plaque psoriasis. These statements deal with future events and involve known and unknown risks, uncertainties and other factors that may cause actual results, performance or achievements to be materially different from the information expressed or implied by these forward-looking statements. Factors that could cause actual results to differ materially include risks and uncertainties such as those relating to the design, implementation and outcomes of Dermira's clinical trials; the outcome of future discussions with regulatory authorities relating to the CIMZIA clinical program; Dermira's dependence on third-party clinical research organizations, manufacturers and suppliers; and Dermira's ability to continue to stay in compliance with applicable laws and regulations. You should refer to the section entitled "Risk Factors" set forth in Dermira's Annual Report on Form 10-K, Dermira's Quarterly Reports on Form 10-Q and other filings Dermira makes with the SEC from time to time for a discussion of important factors that may cause actual results to differ materially from those expressed or implied by Dermira's forward-looking statements. Furthermore, such forward-looking statements speak only as of the date of this press release. Dermira undertakes no obligation to publicly update any forward-looking statements or reasons why actual results might differ, whether as a result of new information, future events or otherwise, except as required by law.

UCB Forward-Looking Statements

This press release contains forward-looking statements based on current plans, estimates and beliefs of management. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including estimates of revenues, operating margins, capital expenditures, cash, other financial information, expected legal, political, regulatory or clinical results and other such estimates and results. By their nature, such forward-looking statements are not guarantees of future performance and are subject to risks, uncertainties and assumptions which could cause actual results to differ materially from those that may be implied by such forward-looking statements contained in this press release. Important factors that could result in such differences include: changes in general economic, business and competitive conditions, the inability to obtain necessary regulatory approvals or to obtain them on acceptable terms, costs associated with research and development, changes in the prospects for products in the pipeline or under development by UCB, effects of future judicial decisions or governmental investigations, product liability claims, challenges to patent protection for products or product candidates, changes in laws or regulations, exchange rate fluctuations, changes or uncertainties in tax laws or the administration of such laws and hiring and retention of its employees. UCB is providing this information as of the date of this press release and expressly disclaims any duty to update any information contained in this press release, either to confirm the actual results or to report a change in its expectations.

There is no guarantee that new product candidates in the pipeline will progress to product approval or that new indications for existing products will be developed and approved. Products or potential products which are the subject of partnerships, joint ventures or licensing collaborations may be subject to differences between the partners. Also, UCB or others could discover safety, side effects or manufacturing problems with its products after they are marketed. Moreover, sales may be impacted by international and domestic trends toward managed care and health care cost containment and the reimbursement policies imposed by third-party payers as well as legislation affecting biopharmaceutical pricing and reimbursement.

Dermira Contacts:

Media:

Erica Jefferson
Senior Director, Head of Corporate Communications
T +1.650.421.7216
media@dermira.com

Investors:

Ian Clements, Ph.D.
Vice President, Investor Relations
T +1.650.422.7753
investor@dermira.com

Robert H. Uhl
Westwicke Partners
Managing Director
T +1.858.356.5932
robert.uhl@westwicke.com

UCB Contacts:

France Nivelles
Global Communications, UCB
T +32.2.559.9178 france.nivelles@ucb.com

Andrea Christopher,
Global Immunology Communications, UCB
T +1.404.483.7329, andrea.levin@ucb.com

Laurent Schots
Media Relations, UCB
T +32.2.559.92.64 Laurent.schots@ucb.com

Antje Witte
Investor Relations, UCB
T +32.2.559.94.14 antje.witte@ucb.com

Isabelle Ghellynck,
Investor Relations, UCB
T +32.2.559.9588, isabelle.ghellynck@ucb.com